Monoidal Rigidity for Free Wreath Products

Pierre Fima and Lorenzo Pittau

Abstract

In this note we observe that any compact quantum group monoidally equivalent, in a nice way, to a free wreath product of a compact quantum group G by the quantum automorphism group of a finite dimensional C*-algebra with a δ-form is actually isomorphic to a free wreath product of G by the quantum automorphism group of another finite dimensional C*-algebra with a δ-form.

arXiv:1604.01236v1 [math.QA] 5 Apr 2016

Introduction

The theory of compact quantum groups, which includes the fundamental constructions of Drinfeld and Jimbo [Dri86, Dri87, Jim85] as q-deformation of the universal enveloping algebras of classical Lie algebras, was introduced by Woronowicz in the eighties [Wor87]. Later, Wang [Wan98] introduced the quantum automorphism group $G_{(B,\psi)}$ of a finite dimensional C*-algebra B with a state ψ and Banica [Ban99] studied its representation category when ψ is a δ-form.

Bichon [Bic04] introduced the construction of free wreath product $G \wr_* S_n^+$ of a compact quantum group G by the permutation quantum group S_n^+ by using an action of S_n^+ on n copies of G. In analogy with the classical case, the free wreath product allows to describe the quantum symmetry group of n copies of a finite graph in terms of the symmetry group of the graph and of S_n^+. The representation category of a free wreath product $G \wr_* S_n^+$, when G is Kac, was computed in [LT14] by Lemeux and Tarrago. The free wreath product $H^+_{(B,\psi)}(G)$ of a compact quantum and a discrete group has been introduced by the second author in the case of $G = \widehat{\Gamma}$, the dual of a discrete group Γ, and the representation category has been computed in [Pit16]. For a general G, the construction and the computation of the representation category has been studied in [FP15].

This note is a continuation of the general study of free wreath product initiated in [FP15]. Since we know from [FP15] the rigid C*-tensor category given by the representation theory of a free wreath product $H^+_{(B,\psi)}(G)$, we will focus now on compact quantum groups which are monoidally equivalent to such a free wreath product. The purpose of this note is to prove the following rigidity result.

Theorem A. *Let G, G_0 be compact quantum groups and (B, ψ) be a finite dimensional C*-algebra with a δ-form ψ. If G is monoidally equivalent to $H^+_{(B,\psi)}(G_0)$ via a* nice enough *unitary tensor functor then there exists a finite dimensional C*-algebra D with a δ-form ψ_D on D such that G is isomorphic to $H^+_{(D,\psi_D)}(G_0)$.*

The precise meaning of *nice enough* will be given in Section 2, just before the statement of Theorem 2.3. When $G_0 = \{e\}$ is the trivial (quantum) group, any unitary tensor functor is nice enough. However, when $G_0 = \{e\}$, a stronger statement than the one of Theorem A is known to hold: it is proved in [Mro15] that any compact quantum group having the same fusion rules as the one of $SO(3)$ is actually isomorphic to the quantum automorphism group of some finite dimensional C*-algebra.

The paper has two sections: Section 1 is a preliminary section in which we introduce the notations of this paper and recall some useful results and Section 2 contains the proof of Theorem A (see Theorem 2.3).

1 Preliminaries

In this paper, we always assume the scalar products on Hilbert spaces to be linear in the first variable. For Hilbert spaces H, K, we denote by $\mathcal{L}(H, K)$ the Banach space of bounded linear maps from H to K and we write $\mathcal{L}(H) := \mathcal{L}(H, H)$. The same symbol $\otimes$ is used to denote the tensor product of Hilbert spaces or the minimal tensor product of C*-algebras.

Given a unital C*-algebra A and unitaries $u \in \mathcal{L}(H) \otimes A$ and $v \in \mathcal{L}(K) \otimes A$, for H, K Hilbert spaces, we define $\mathrm{Mor}(u, v) := \{T \in \mathcal{L}(H, K) \,:\, (T \otimes 1)u = v(T \otimes 1)\}$. We also use the notation

$$u \otimes v := u_{13} v_{23} \in \mathcal{L}(H \otimes K) \otimes A,$$

where $v_{23} := \mathrm{id}_H \otimes v$ and $u_{13} := (\sigma \otimes \mathrm{id}_A)(\mathrm{id}_K \otimes u)$ and $\sigma \,:\, \mathcal{L}(K) \otimes \mathcal{L}(H) \to \mathcal{L}(H) \otimes \mathcal{L}(K)$ is the flip automorphism.

For G a compact quantum group in the sense of [Wor98] we denote by $C(G)$ the maximal C*-algebra of G i.e. the enveloping C*-algebra of the $*$-algebra given by the linear span of the coefficients of the irreducible representations of G. We will also denote by ε the trivial representation of G, $\mathrm{Irr}(G)$ the set of equivalence classes of irreducible representations of G and by $\mathcal{R}(G)$ the rigid C*-tensor category of finite dimensional unitary representations of G. We refer to the chapter 2 of the book [NT13] for all the details concerning rigid C*-tensor categories and monoidal equivalence.

Let B be a finite dimensional C*-algebra and ψ a faithful state on B. Then B is a Hilbert space with the scalar product defined by $\langle a, b\rangle := \psi(b^* a)$. Let $m_B \,:\, B \otimes B \to B$ be the multiplication on B and $\eta_B \,:\, \mathbb{C} \to B$ be the unit. We call ψ a *δ-form* when $m_B m_B^* = \delta \mathrm{id}_B$, where $m_B^* \,:\, B \to B \otimes B$ is the adjoint of m_B with respect to the scalar product on B defined above.

Let G_0 be a compact quantum group and for $\alpha \in \mathrm{Irr}(G_0)$, we choose a representative u^α acting on the Hilbert space H_α. The free wreath product of G_0 by the quantum automorphism group $G_{(B,\psi)}$ of (B, ψ), denoted by $H^+_{(B,\psi)}(G_0)$, is the compact quantum group defined in [FP15] as follows. The full C*-algebra $C(H^+_{(B,\psi)}(G_0))$ of this quantum group is the universal unital C*-algebra generated by the coefficients of matrices $a(\alpha) \in \mathcal{L}(B \otimes H_\alpha) \otimes C(H^+_{(B,\psi)}(G_0))$, for $\alpha \in \mathrm{Irr}(G_0)$, such that:

- $a(\alpha)$ is unitary for all $\alpha \in \mathrm{Irr}(G_0)$,
- $(m_B \otimes S)\Sigma_{23} \in \mathrm{Mor}(a(\alpha) \otimes a(\beta), a(\gamma))$ for all $\alpha, \beta, \gamma \in \mathrm{Irr}(G_0)$ and all $S \in \mathrm{Mor}(\alpha \otimes \beta, \gamma)$,
- $\eta_B \in \mathrm{Mor}(\varepsilon, a(\varepsilon))$.

Moreover, there exists a unique unital $*$-homomorphism $\Delta \,:\, C(H^+_{(B,\psi)}(G_0)) \to C(H^+_{(B,\psi)}(G_0)) \otimes C(H^+_{(B,\psi)}(G_0))$ for which the elements $a(\alpha)$ are representations for all $\alpha \in \mathrm{Irr}(G_0)$. The pair $(C(H^+_{(B,\psi)}(G_0)), \Delta)$ is a compact quantum group, called the free wreath product.

When ψ is a δ-form, the representation theory of $H^+_{(B,\psi)}(G_0)$ is totally understood [FP15]. In particular, the dimension of the spaces $\mathrm{Mor}(\varepsilon, a(\alpha_1) \otimes \ldots \otimes a(\alpha_n))$ only depends on G_0 and $\alpha_1, \ldots, \alpha_n$ and not on (B, ψ).

2 Proof of Theorem A

Suppose that G and G_0 are compact quantum groups and G is unitary monoidally equivalent to $H^+_{(B,\psi)}(G_0)$, where ψ is a δ-form. This means (see [NT13]) that there exist unitary tensor functors $F \,:\, \mathcal{R}(H^+_{(B,\psi)}(G_0)) \to \mathcal{R}(G)$, $H \,:\, \mathcal{R}(G) \to \mathcal{R}(H^+_{(B,\psi)}(G_0))$ such that FH and HF are naturally monoidally unitarily isomorphic to the identity functors, meaning that the natural isomorphisms $FH \simeq \iota$ and $HF \simeq \iota$ are both implemented by unitaries.

For $\alpha \in \mathrm{Irr}(G_0)$, we choose a representative u^α acting on the Hilbert space H_α. Recall that the representation $a(\alpha)$ is acting on $B \otimes H_\alpha$. We define the unitary representation $v_\alpha := F(a(\alpha))$ of G acting on the finite dimensional Hilbert space K_α. For all u, v finite dimensional unitary representations of $H^+_{(B,\psi)}(G_0)$, we denote by the same symbol $F_2 \in \mathrm{Mor}(F(u) \otimes F(v), F(u \otimes v))$ the unitary given by the definition of the unitary tensor functor F. Since the category is strict we have:

$$F_2 \circ (F_2 \otimes \mathrm{id}) = F_2 \circ (\mathrm{id} \otimes F_2) \in \mathrm{Mor}(F(u) \otimes F(v) \otimes F(w), F(u \otimes v \otimes w)) \text{ for all } u, v, w. \tag{2.1}$$

Moreover, the naturally of F_2 means that, for all u, v, u', v' finite dimensional unitary representations we have:

$$F(f \otimes g) \circ F_2 = F_2 \circ (F(f) \otimes F(g)) \in \mathrm{Mor}(F(u) \otimes F(v), F(u' \otimes v')) \quad \text{for all } f \in \mathrm{Mor}(u, u'), g \in \mathrm{Mor}(v, v'). \tag{2.2}$$

Finally, we note that, since F is unitary, we may and will assume that

$$F(\varepsilon) = \varepsilon \quad \text{and} \quad F(\mathrm{id}) = \mathrm{id} \text{ for all } \mathrm{id} \in \mathrm{Mor}(u, u). \tag{2.3}$$

Define the finite dimensional Hilbert space $D = K_\varepsilon$. Since $m_B \in \mathrm{Mor}(a(\varepsilon)\otimes a(\varepsilon), a(\varepsilon))$ and $\eta_B \in \mathrm{Mor}(\varepsilon, a(\varepsilon))$, we may define the linear maps

$$m_D : D \otimes D \to D,\ m_D := F(m_B)F_2 \quad \text{and} \quad \eta_D : \mathbb{C} \to D,\ \eta_D := F(\eta_B).$$

Proposition 2.1. *(D, m_D, η_D) is a unital algebra and $m_D m_D^* = \delta \mathrm{id}_D$.*

Proof. Using Equations (2.1), (2.2), (2.3) and the associativity of m_B we get:

$$\begin{aligned} m_D \circ (m_D \otimes \mathrm{id}) &= F(m_B)F_2 \circ (F(m_B)F_2 \otimes \mathrm{id}) = F(m_B) \circ F_2 \circ (F(m_B) \otimes \mathrm{id}) \circ (F_2 \otimes 1) \\ &= F(m_B) \circ F(m_B \otimes \mathrm{id}) \circ F_2 \circ (F_2 \otimes 1) = F(m_B \circ (m_B \otimes \mathrm{id})) \circ F_2 \circ (\mathrm{id} \otimes F_2) \\ &= F(m_B \circ (\mathrm{id} \otimes m_B)) \circ F_2 \circ (\mathrm{id} \otimes F_2) = F(m_B) \circ F(\mathrm{id} \otimes m_B) \circ F_2 \circ (\mathrm{id} \otimes F_2) \\ &= F(m_B) \circ F_2 \circ (\mathrm{id} \otimes F(m_B)) \circ (\mathrm{id} \otimes F_2) = F(m_B)F_2 \circ (\mathrm{id} \otimes F(m_B)F_2) \\ &= m_D \circ (\mathrm{id} \otimes m_D). \end{aligned}$$

Hence, m_D is associative. Moreover, using Equations (2.2), (2.3) and the the fact that η_B is the unit of (B, m_B), which means that $m_B(\eta_B \otimes \mathrm{id}_B) = \mathrm{id}_B = m_B(\mathrm{id}_B \otimes \eta_B)$, we find:

$m_D(\eta_D \otimes \mathrm{id}_D) = F(m_B)F_2(F(\eta_B)\otimes \mathrm{id}_D) = F(m_B)F_2F_2^*F(\eta_B \otimes \mathrm{id}_B) = F(m_B(\eta_B \otimes \mathrm{id}_B)) = F(\mathrm{id}_B) = \mathrm{id}_D$ and,

$m_D(\mathrm{id}_D \otimes \eta_D) = F(m_B)F_2(\mathrm{id}_D \otimes F(\eta_B)) = F(m_B)F_2F_2^*F(\mathrm{id}_B \otimes \eta_B) = F(m_B(\mathrm{id}_B \otimes \eta_B)) = F(\mathrm{id}_B) = \mathrm{id}_D$.

Finally, $m_D m_D^* = F(m_B) \circ F_2 \circ F_2^* \circ F(m_B)^* = F(m_B m_B^*) = \delta F(\mathrm{id}_B) = \delta \mathrm{id}_D$. □

We are now ready to turn D into a $*$-algebra. Define $t = m_D^* \circ \eta_D = F_2^* F(m_B^* \eta_B) \in \mathcal{L}(\mathbb{C}, D \otimes D)$. We may and will view $t \in D \otimes D$. Define the anti-linear map $S_D : D \to D$ by $S_D(x) = (x^* \otimes \mathrm{id})(t)$. Denote by $L_x \in \mathcal{L}(D)$ the bounded operator given by left multiplication by $x \in D$ and write $1_D := \eta_D(1) \in D$.

Proposition 2.2. *S_D is an involution on the unital algebra D. Moreover, equipped with this involution and the norm defined by $\|x\|_D := \|L_x\|$, for $x \in D$, D is a unital C*-algebra and the following holds.*

1. *$(L_x)^* = L_{S_D(x)}$ for all $x \in D$.*
2. *$\psi_D : D \to \mathbb{C}$, $x \mapsto \langle x, 1_D\rangle$ is a faithful state on D satisfying $\psi_D(S_D(y)x) = \langle x, y\rangle$ for all $x, y \in D$.*
3. *ψ_D is a δ-form on D.*

Proof. (1) Using diagrams computations one can easily check that:

$$(\mathrm{id} \otimes m_B) \circ (m_B^* \circ \eta_B \otimes \mathrm{id}) = m_B^*.$$

Applying the functor F we find $(\mathrm{id} \otimes m_D) \circ (m_D^* \circ \eta_D \otimes \mathrm{id}) = m_D^*$. Hence, for all $x, y, z \in D$, one has

$$\begin{aligned} \langle L_{S_D(x)}(y), z\rangle &= \langle m_D(S_D(x) \otimes y), z\rangle = \langle m_D((x^* \otimes \mathrm{id}) \circ m_D^* \circ \eta_D \otimes y), z\rangle \\ &= \langle (x^* \otimes \mathrm{id})(\mathrm{id} \otimes m_D) \circ (m_D^* \circ \eta_D \otimes \mathrm{id})(y), z\rangle = \langle (x^* \otimes \mathrm{id})(m_D^*(y)), z\rangle \\ &= \langle m_D^*(y), x \otimes z\rangle = \langle y, m_D(x \otimes z)\rangle = \langle y, L_x(z)\rangle = \langle (L_x)^*(y), z\rangle \end{aligned}$$

This concludes the proof of 1 and shows that S_D is anti-multiplicative since it implies that, for all $x, y \in D$,

$$S_D(xy) = (L_{xy})^*(1_D) = (L_y)^*(L_x)^*(1_D) = L_{S_D(y)}(S_D(x)) = S_D(y)S_D(x).$$

Assertion (1) also shows that S_D is involutive since it implies that $L_{S_D^2(x)} = (L_{S_D(x)})^* = ((L_x)^*)^* = L_x$ hence, $S_D^2(x) = x$. It follows that S_D turns (D, m_D, η_D) into an involutive unital algebra. It is moreover

clear that $\|x\|_D := \|L_x\|_{\mathcal{L}(D)}$, for $x \in D$, defines a norm on D which satisfies the C*-condition since we have $\|S_D(x)x\|_D = \|L_{S_D(x)x}\| = \|L_{S_D(x)}L_x\| = \|L_x^*L_x\| = \|L_x\|^2 = \|x\|_D$ for all $x \in D$.

(2) One has $\psi_D(S_D(y)x) = \langle S_D(y)x, 1_D\rangle = \langle L_{S_D(y)}(x), 1_D\rangle = \langle (L_y)^*(x), 1_D\rangle = \langle x, L_y(1_D)\rangle = \langle x, y\rangle$. It follows directly from this relation and the non degeneracy of the scalar product on D that ψ_D is faithful. It also follows that the scalar product induced by ψ_D on D is the same as the initial scalar product on D. Hence, taking the adjoint with the ψ_D-scalar product we have, from Proposition 2.1, $m_D m_D^* = \delta \mathrm{id}_D$. □

We assume from now that the functor F is *nice enough* meaning that for any $\alpha \in \mathrm{Irr}(G_0)$, there exists a unitary $V_\alpha \in \mathcal{L}(K_\alpha, K_\varepsilon \otimes H_\alpha)$, with $V_\varepsilon = \mathrm{id}_D$, and such that the following diagram is commutative for all $\alpha, \beta, \gamma \in \mathrm{Irr}(G_0)$ and all $S \in \mathrm{Mor}(\alpha \otimes \beta, \gamma)$:

$$\begin{array}{ccc} (K_\varepsilon \otimes H_\alpha) \otimes (K_\varepsilon \otimes H_\beta) & \overset{(m_D \otimes S)\Sigma_{23}}{\longrightarrow} & K_\varepsilon \otimes H_\gamma \\ \uparrow V_\alpha \otimes V_\beta & & \uparrow V_\gamma \\ K_\alpha \otimes K_\beta & \overset{F((m_B \otimes S)\Sigma_{23})F_2}{\longrightarrow} & K_\gamma \end{array}$$

Observe that any unitary tensor functor is nice enough if G_0 is the trivial quantum group. We are now ready to state our main result. Let us denote by $\widetilde{a}(\alpha) \in \mathcal{L}(D \otimes H_\alpha) \otimes C(H^+_{(D,\psi_D)}(G_0))$ the canonical generators.

Theorem 2.3. *There exists a unique $*$-isomorphism $\pi : C(H^+_{(D,\psi_D)}(G_0)) \to C(G)$ such that*

$$(\mathrm{id} \otimes \pi)(\widetilde{a}(\alpha)) = (V_\alpha \otimes 1)v_\alpha(V_\alpha^* \otimes 1) \in \mathcal{L}(D \otimes H_\alpha) \otimes C(G) \quad \textit{for all } \alpha \in \mathrm{Irr}(G_0).$$

Moreover, π intertwines the comultiplications.

Proof. **Step 1.** *There exists a surjective unital $*$-homomorphism $\pi : C(H^+_{(D,\psi_D)}(G_0)) \to C(G)$ such that $(\mathrm{id} \otimes \pi)(\widetilde{a}(\alpha)) = (V_\alpha \otimes 1)v_\alpha(V_\alpha^* \otimes 1) \in \mathcal{L}(D \otimes H_\alpha) \otimes C(G)$ for all $\alpha \in \mathrm{Irr}(G_0)$.*

From the universal property of the C*-algebra $C(H^+_{(D,\psi_D)}(G_0))$ it suffices to check that the unitary representations $u_\alpha := (V_\alpha \otimes 1)v_\alpha(V_\alpha^* \otimes 1)$ of G on the Hilbert space $D \otimes H_\alpha$ satisfy the following relations:

1. $(m_D \otimes S)\Sigma_{23} \in \mathrm{Mor}(u_\alpha \otimes u_\beta, u_\gamma)$ for all $\alpha, \beta, \gamma \in \mathrm{Irr}(G_0)$ and all $S \in \mathrm{Mor}(\alpha \otimes \beta, \gamma)$,
2. $\eta_D \in \mathrm{Mor}(\varepsilon, u_\varepsilon)$).

(1) Since F is nice enough we have, for all $\alpha, \beta, \gamma \in \mathrm{Irr}(G_0)$ and all $S \in \mathrm{Mor}(\alpha \otimes \beta, \gamma)$,

$$V_\gamma^*(m_D \otimes S)\Sigma_{23}(V_\alpha \otimes V_\beta) = F((m_B \otimes S)\Sigma_{23})F_2 \in \mathrm{Mor}(v_\alpha \otimes v_\beta, v_\gamma).$$

It follows that $(((m_D \otimes S)\Sigma_{23}) \otimes 1)(u_\alpha \otimes u_\beta)$ is equal to:

$$\begin{aligned} &(((m_D \otimes S)\Sigma_{23}) \otimes 1)(V_\alpha \otimes 1 \otimes 1)(v_\alpha)_{13}(V_\alpha^* \otimes 1 \otimes 1)(1 \otimes V_\beta \otimes 1)(v_\beta)_{23}(1 \otimes V_\beta^* \otimes 1) \\ &= ((((m_D \otimes S)\Sigma_{23})(V_\alpha \otimes V_\beta)) \otimes 1)(v_\alpha \otimes v_\beta)(V_\alpha^* \otimes V_\beta^* \otimes 1) \\ &= (V_\gamma \otimes 1)v_\gamma(V_\gamma^*((m_D \otimes S)\Sigma_{23})(V_\alpha \otimes V_\beta)(V_\alpha^* \otimes V_\beta^*) \otimes 1) = u_\gamma((m_D \otimes S)\Sigma_{23} \otimes 1). \end{aligned}$$

(2). It is obvious since $\eta_B \in \mathrm{Mor}(\varepsilon, a(\varepsilon))$ implies that $\eta_D = F(\eta_B) \in \mathrm{Mor}(\varepsilon, v_\varepsilon)$ and $u_\varepsilon = v_\varepsilon$.

Note that π automatically intertwines the comultiplications and π is surjective since F is essentially surjective.

Step 2. *π is an isomorphism.*

It suffices to check that π intertwines the Haar measures h on $C(H^+_{(D,\psi_D)}(G_0))$ and h_G on $C(G)$. Since the linear span of the coefficients of representations of the form $\widetilde{a}(\alpha_1) \otimes \ldots \otimes \widetilde{a}(\alpha_n)$, for $\alpha_1, \ldots, \alpha_n \in \mathrm{Irr}(G_0)$ and $n \geq 1$, is dense in $C(H^+_{(D,\psi_D)}(G_0))$, it suffices to check that, for all $n \geq 1$, $\alpha_1, \ldots, \alpha_n \in \mathrm{Irr}(G_0)$ one has

$$(\mathrm{id} \otimes h_G)(u_{\alpha_1} \otimes \ldots \otimes u_{\alpha_n}) = (\mathrm{id} \otimes h)(\widetilde{a}(\alpha_1) \otimes \ldots \otimes \widetilde{a}(\alpha_n)),$$

which is equivalent to $\dim(\mathrm{Mor}(\varepsilon, u_{\alpha_1}\otimes\ldots\otimes u_{\alpha_n})) = \dim(\mathrm{Mor}(\varepsilon, \widetilde{a}(\alpha_1)\otimes\ldots\otimes\widetilde{a}(\alpha_n))$. Since $u_\alpha \simeq v_\alpha = F(a(\alpha))$ for all $\alpha \in \mathrm{Irr}(G_0)$ and F is a monoidal equivalence, the left hand side is equal to

$$\dim(\mathrm{Mor}(\varepsilon, v_{\alpha_1} \otimes \ldots \otimes v_{\alpha_n})) = \dim(\mathrm{Mor}(\varepsilon, a(\alpha_1) \otimes \ldots \otimes a(\alpha_n)).$$

Moreover since, ψ_D is a δ-form we also know from [FP15, Theorem 3.5] an explicit formula for the number $\dim(\mathrm{Mor}(\varepsilon, a(\alpha_1) \otimes \ldots \otimes a(\alpha_n))$ which only depends on G_0 and not on (B, ψ) or (D, ψ_D) and we have

$$\dim(\mathrm{Mor}(\varepsilon, a(\alpha_1) \otimes \ldots \otimes a(\alpha_n)) = \dim(\mathrm{Mor}(\varepsilon, \widetilde{a}(\alpha_1) \otimes \ldots \otimes \widetilde{a}(\alpha_n)).$$

□

References

[Ban99] Teodor Banica. Symmetries of a generic coaction. *Math. Ann.*, 314(4):763–780, 1999.

[Bic04] Julien Bichon. Free wreath product by the quantum permutation group. *Algebr. Represent. Theory*, 7(4):343–362, 2004.

[Dri86] V. G. Drinfel′d. Quantum groups. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 155(Differentsialnaya Geometriya, Gruppy Li i Mekh. VIII):18–49, 193, 1986.

[Dri87] V. G. Drinfel′d. Quantum groups. In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986)*, pages 798–820. Amer. Math. Soc., Providence, RI, 1987.

[FP15] Pierre Fima and Lorenzo Pittau. The free wreath product of a compact quantum group by a quantum automorphism group. *preprint arXiv:1507.06107*, 2015.

[Jim85] Michio Jimbo. A q-difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation. *Lett. Math. Phys.*, 10(1):63–69, 1985.

[LT14] François Lemeux and Pierre Tarrago. Free wreath product quantum groups: the monoidal category, approximation properties and free probability. *preprint arXiv:1411.4124*, 2014.

[Mro15] Colin Mrozinski. Quantum automorphism groups and so(3)-deformations. *J. Pure Appl. Algebra*, 219(1):1–32, 2015.

[NT13] Sergey Neshveyev and Lars Tuset. *Compact quantum groups and their representation categories*, volume 20 of *Cours Spécialisés [Specialized Courses]*. Société Mathématique de France, Paris, 2013.

[Pit16] Lorenzo Pittau. The free wreath product of a discrete group by a quantum automorphism group. *Proc. Amer. Math. Soc.*, 144(5):1985–21, 2016.

[Wan98] Shuzhou Wang. Quantum symmetry groups of finite spaces. *Comm. Math. Phys.*, 195(1):195–211, 1998.

[Wor87] S. L. Woronowicz. Compact matrix pseudogroups. *Comm. Math. Phys.*, 111(4):613–665, 1987.

[Wor98] S. L. Woronowicz. Compact quantum groups. In *Symétries quantiques (Les Houches, 1995)*, pages 845–884. North-Holland, Amsterdam, 1998.

Pierre FIMA
Univ Paris Diderot, Sorbonne Paris Cité, IMJ-PRG, UMR 7586, F-75013, Paris, France
Sorbonne Universités, UPMC Paris 06, UMR 7586, IMJ-PRG, F-75005, Paris, France
CNRS, UMR 7586, IMJ-PRG, F-75005, Paris, France
E-mail address: pierre.fima@imj-prg.fr

Lorenzo PITTAU
E-mail address: lorenzopittau@gmail.com